

# Improved Half-life Measurement of $^{224}\text{Pa}$ and its $^{209}\text{Bi}(^{18}\text{O},3n)^{224}\text{Pa}$ Production Cross Section

Philip A. Wilk, Ken E. Gregorich, Melissa B. Hendricks, Mike R. Lane, Diana M. Lee, Chris A. McGrath,  
Dawn A. Shaughnessy, Dan A. Strellis, Eric R. Sylwester, and Darleane C. Hoffman

Two half-lives for  $^{224}\text{Pa}$  have been reported previously. However, the error bars of these two measurements,  $600\pm 50$  ms measured for  $^{224}\text{Pa}$  from the  $^{232}\text{Th}(p,9n)^{224}\text{Pa}$  reaction<sup>1</sup> and  $950\pm 150$  ms measured for  $^{224}\text{Pa}$  from the  $^{205}\text{Tl}(^{22}\text{Ne},3n)^{224}\text{Pa}$  reaction<sup>2</sup>, do not overlap. The former measurement's low value could be due to misinterpretation of the very complex alpha spectrum that was obtained. The disparity could lie in the misidentification of a  $^{223}\text{Th}$  ( $t_{1/2}=650$  ms,  $a=7.324$  MeV) decay peak produced by the  $^{232}\text{Th}(p,p9n)^{224}\text{Pa}$  reaction.

In our experiment  $^{224}\text{Pa}$  was produced via the  $^{209}\text{Bi}(^{18}\text{O}, 3n)^{224}\text{Pa}$  reaction which has been previously reported<sup>3</sup>, but no production cross section or additional half-life measurement was given. The Lawrence Berkeley National Laboratory 88-Inch Cyclotron provided a 20 particle-nA beam of 111 MeV (laboratory frame)  $^{18}\text{O}^{5+}$ . The target system has been described in detail previously<sup>4</sup>. The energy calibration was performed on-line using known alpha decay energies from the following nuclides:  $^{211}\text{Bi}$ ,  $^{214}\text{Ra}$ ,  $^{211}\text{Po}$ ,  $^{212}\text{Ac}$ ,  $^{216}\text{Fr}$ ,

The odd-odd nucleus  $^{224}\text{Pa}$  is not expected to have a significant beta-decay branch and appears to alpha decay mainly to a single nuclear level in  $^{220}\text{Ac}$  with an alpha particle energy of 7.49 MeV<sup>2,5</sup>. Unfortunately, while alpha particles with this energy are identifiable in our alpha-energy spectrum, they are largely obscured by the  $^{211}\text{Po}$  transfer product which decays via a 7.45 MeV alpha. Therefore the  $^{224}\text{Pa}$  decay was identified by observing its alpha-decay daughters  $^{220}\text{Ac}$  ( $t_{1/2}=26$  ms;  $\alpha=7.610, 7.680, 7.790, 7.850$  MeV)<sup>5</sup> and  $^{216}\text{Fr}$  ( $t_{1/2}=0.7$  ms;  $\alpha=9.01$  MeV)<sup>5</sup>. The  $^{224}\text{Pa}$  daughter events were observed with 6 pairs of opposing detectors<sup>6</sup> within 2  $\mu\text{s}$  of each other. By fitting the resultant decay curve with two components, the half-life of  $^{224}\text{Pa}$  was determined to be  $850\pm 20$  ms

(see Fig. 1). From these data, we also determined the production cross section to be  $0.5\pm 0.1$  mb, after taking into account the detector efficiency, capillary transport efficiency, and capillary transport time.

The half-life for  $^{224}\text{Pa}$  as determined by this experiment is consistent with, but much more precise than the value of  $950\pm 150$  ms reported by Borggreen *et al.*<sup>2</sup>. Our measured production cross-section is consistent with the value of 0.35 mb calculated with the SPIT code<sup>7</sup>.

## Footnotes and References

1. P. A. Tove, Arkiv Fysik. **13**, 549 (1958).
2. J. Borggreen, K. Valli, E. K. Hyde. Phys. Rev. C **2**, 1841 (1970).
3. M. Fukuda et al., RIKEN Accel. Prog. Rep. **20**, 14 (1986).
4. D. Lee *et al.*, Phys. Rev. C **25**, 286 (1982).
5. R. B. Firestone, Table of Isotopes, 8th ed. (Wiley-Interscience, New York, 1996).
6. D. C. Hoffman *et al.*, Phys. Rev. C **41**, 631 (1990).
7. T. Sikkeland, Ark. Fys. **36**, 539 (1967); J. Alonso, Gmelin Handbook of Inorg. Chem., **7b**, 28 (1974).




Fig. 1. Decay of  $^{224}\text{Pa}$